

# P506

## Problem Set 1

(Due: September 11, 2008)

1) Jackson, problem 1.1.

Use Gauss's theorem [and (1.21) if necessary] to prove the following:

- (a) Any excess charge placed on a conductor must lie entirely on its surface. (A conductor by definition contains charges capable of moving freely under the action of applied electric fields.)
- (b) A closed, hollow conductor shields its interior from fields due to charges outside, but does not shield its exterior from the fields due to charges placed inside it.
- (c) The electric field at the surface of a conductor is normal to the surface and has a magnitude  $\sigma/\epsilon_0$ , where  $\sigma$  is the charge density per unit area on the surface.

2) Jackson, problem 1.3.

Using the Dirac delta function in the appropriate coordinates, express the following charge distributions as three-dimensional charge densities  $\rho(\mathbf{x})$ .

- (a) In spherical coordinates, a charge  $Q$  uniformly distributed over a spherical shell of radius  $R$ .
- (b) In cylindrical coordinates, a charge  $\lambda$  per unit length uniformly distributed over a cylindrical surface of radius  $b$ .
- (c) In cylindrical coordinates, a charge  $Q$  spread uniformly over a flat circular disc of negligible thickness and radius  $R$ .
- (d) The same as part (c), but using spherical coordinates.

3) Jackson, problem 1.4.

Each of three charged spheres of radius  $a$ , one conducting, one having a uniform charge density within its volume, and one having a spherically symmetric charge density that varies radially as  $r^n$  ( $n > -3$ ), has a total charge  $Q$ . Use Gauss's theorem to obtain the electric fields both inside and outside each sphere. Sketch the behavior of the fields as a function of radius for the first two spheres, and for the third with  $n = -2, +2$ .

4) Jackson, problem 1.5.

The time-averaged potential of a neutral hydrogen atom is given by

$$\Phi = \frac{q}{4\pi\epsilon_0} \frac{e^{-\alpha r}}{r} \left( 1 + \frac{\alpha r}{2} \right)$$

where  $q$  is the magnitude of the electronic charge, and  $\alpha^{-1} = a_0/2$ ,  $a_0$  being the Bohr radius. Find the distribution of charge (both continuous and discrete) that will give this potential and interpret your result physically.